Joseph Benzel

1) Identification Technology Program (ITP) / Colorado State University, USDA-APHIS-PPQ-Science & Technology (S&T), 2301 Research Boulevard, Suite 108, Fort Collins, Colorado 80526 U.S.A. (Email: itp@USDA.gov)

Version 3 18 June 2015 This CAPS (Cooperative Agricultural Pest Survey) screening aid produced for and distributed by: USDA-APHIS-PPQ National Identification Services (NIS) This and other identification resources are available at: http://caps.ceris.purdue.edu/taxonomic_services



The redbay ambrosia beetle, *Xyleborus glabratus* Eichhoff (Fig. 1) is an economically important pest native to India, Japan, and southeast Asia. The primary hosts of this species are trees in the laurel family (Lauracea) such as redbay (*Persia borbonia*), avocado (*P. americanum*) and Asian spicebush (*Lindera latifolia*). The beetles and their larvae cultivate and feed on the symbiotic fungus *Raffelea* sp. which causes laurel wilt disease and can kill healthy trees (Figs. 2-4).

Xyleborus glabratus is a member of the Curculionidae (subfamily Scolytinae) which is comprised of weevils and bark beetles. Members of this family are highly variable but almost all species share a distinct club on the end of their antennae consisting of three segments. The subfamily Scolytinae, to which *Xyleborus* belongs, consists of the bark beetles. In general, members of Scolytinae are small (<10mm long) pill shaped beetles of a reddish brown or black, or tan color. Some authors consider Scolytinae to be a distinct family (Scolytidae).

The tribe Xyleborini is a large and widespread group of genera of the Scolytinae. The tribe is most notable for having dwarfed haploid males which never leave the parental galleries (Fig. 5-6) and numerous small socketed spines on the tibia of females. The genus *Xyleborus* contains more than 500 species, seventeen of which are native to North America. The genus has a worldwide distribution, inhabiting every continent except Antarctica. The genus *Xyleborus* is distinguished by an obliquely truncate club, elongate pronotum, and flat scutellum. *Xyleborus glabratus* was detected in 2002 in Georgia and has since spread to South Carolina in 2004, Florida in 2009, and by 2011 in North Carolina, Alabama and possibly Mississippi. Non-targets often captured during *X. glabratus* surveys include other members of the genus such as *X. celsus*, *X. dispar*, and *X. ferrugenous*. A quality, high powered microscope is required to examine the characters necessary to identify these beetles.

This aid is designed to assist in the sorting and screening of *X. glabratus* suspect adults collected in CAPS Lindgren funnel traps in the continental United States. It covers basic Sorting of traps, First Level, and Second Level screening, all based on morphological characters. Basic knowledge of Coleoptera morphology is necessary to screen for *X. glabratus* suspects.



Fig. 1: *Xyleborus glabratus* on tree (photo by R. Scott Cameron, Advanced Forest Protection, Inc., Bugwood.org).



Fig. 2: *Xyleborus glabratus* galleries (photo by James Johnson, Georgia Forestry Commission, Bugwood.org).

Sorting

Xyleborus glabratus Eichhoff

Xyleborus glabratus survey Lindgren funnel traps should be sorted initially for the presence of beetles of the appropriate size color and shape.

- 1. Beetles are between 2 mm (0.1 inches) and 3 mm (0.2 inches) in length.
- 2. Beetles are pill-like in shape.
- 3. Beetles are black, reddish-brown, or tan colored.

Beetles meeting these requirements should be forwarded to Level 1 Screening (Page 3).



Fig. 3 and 4: Redbay trees killed by laurel wilt. This disease is caused by the fungus *Raffelea* sp. which is spread and cultivated by *X. glabratus*. After the introduction and spread of *X. glabratus* through The Southeastern United States, laurel wilt decimated the areas redbay populations with some areas seeing 90% mortality (photos by Albert (Bud) Mayfield, USDA Forest Service, Bugwood.org).





Fig. 5 and 6: Male (top) and female (bottom) *Xyleborus* sp. All members of the tribe Xyleborini show strong sexual dimorphism with haploid, wingless males that never leave the parental gallery. Females need not mate to produce male offspring and will mate with their own sons. This reproductive strategy allows the beetles to rapidly colonize any suitable area and accounts for the large number of species within the genus (photos by. Maja Jurc, University of Ljubljana, Bugwood.org).

Level 1 Screening

Suspect adults should be pointed and properly labeled. Level 1 Screening is based on characteristics of the antennae, general dorsal surface, and tibia. It is designed to seperate *Xyleborus* from related genera such as *lps* (Figs. 9-10). Specimens with these traits should be forwarded to Level 2 Screening.

Antennae

Scolytids have relatively stout, geniculate, clubbed antennae. The clubs are made up of three antennomers and can be solid, annulated, or occasionally lamellate. The scape will always be well developed (Fig. 6). In *Xyleborus* the antennae have five segmented funicles and obliquely truncate clubs.

General Dorsal Surface

Beetles in the tribe Xyleborini have the basal margins of their elytra unarmed and forming a straight line across the body (Fig. 7). The scutellum of tribe members is usually large and flat and the head is usually concealed at least partially by an enlarged pronotum.

Tibia

The mesothorasic and metathorasic tibia of *Xyleborus* are armed by a series of small teeth on the lateral apical angle (Fig. 8).



Redbay Ambrosia Beetle

Xyleborus glabratus Eichhoff



Fig. 6: Antenna of *Xyleborus* spp. Note the long scape and large three part club.



Fig. 7: Base of elytra and scutellum of *Xyleborus* spp. Members of the Scolytinae have a broad scutellum and the base of the elytra is unarmed.



Fig. 8: *Xyleborus* spp. metathoracic tibia Note the row of small spines on the lateral apical angle (circled).

Figs. 9 and 10 (left): comparison of *Xyleborus glabratus* (far left) and *Ips* spp. (left). Though superficially similar, the Xyleborini and Ipini tribes can be differentiated by the excavated and spined declivity in *Ips*.

Level 2 Screening

Level 2 screening is designed to separate suspect *X. glabratus* (Fig. 15) from other members of *Xyleborus* (Figs. 16-20). Screening is based mostly on general features and the elytral declivity. There are a number of remarkably similar species in this genus, so care should be taken in screening at this level.

General Features

Xyleborus glabratus is a relatively slender species with a length to width ratio of 2.6 or greater (Fig. 15). *Xyleborus sayi* and some other species are more stout (Fig. 18) and are significantly larger than *X. glabratus,* with lengths of around 3-4 mm compared to the 1-2 mm of *X. glabratus.*

Elytral Declivity

The elytral declivity of *X. glabratus* (Figs 21-22) is abruptly flattened or sometimes weakly convex. In addion the declivity ibears a weak but distinct carinate ridge on the lower portion of the lateral margin (circled in Fig. 13) not found in native *Xyleborus* (Fig. 14). The declivity itself is armed with two distinct tubercles in the middle of the first and third interstrae (Figs. 11-12). The declivities of native *Xyleborus* species (Figs. 23-28) are strongly convex or excavated, lack a distinct carina on the lower lateral margin, and have different arrangments of tubercles. Because of the difficulty in identifying these traits, any *Xyleborus* with one or more of the features described above should be considered a viable *X. glabratus* suspect.





Figs. 11-12: Postrolateral and lateral views of the declivity of *X. glabratus*. Note the two small tubercles on the first and third interstria (circled).



Figs. 13 and 14: Lateral views of *X. glabratus* (left) and *X. pubescens* (right). Note the carina on the lower lateral margin of the *X. glabratus* declivity (circled).

Level 2 Non-targets

Redbay Ambrosia Beetle Xyleborus glabratus Eichhoff



Level 2 Non-targets

Redbay Ambrosia Beetle

Xyleborus glabratus Eichhoff



Fig. 21: *Xyleborus glabratus* declivity, posterolateral view (target).



Fig. 22 *Xyleborus glabratus* declivity, lateral view (target).



Fig. 24: *Xyleborus celsus* declivity, lateral view.



Fig. 27: *Xyleborus sayi* declivity, posterolateral view.



Fig. 25: *Xyleborus xylegraphus* declivity, posterolateral view.



Fig. 28: *Xyleborus pubescens* declivity, posterolateral view.



Fig. 23: *Xyleborus celsus* declivity, posterolateral view.



Fig. 26: *Xyleborus ferrugineus* declivity, posterolateral view.

Suspect *X. glabratus* specimens (scolytids with unarmed elytral bases, meso and metatibia armed with small spines, obliquely truncate antennae, and flattened or weakly convex declivities armed a carina on the lower lateral margin and tubercles on the first and third interstria) should be sent forward for identification. Specimens must be labeled and carefully packed to avoid damage during shipping.

Key to Sort and Screen X. glabratus Suspects in the United States

tan.....Not X. glabratus

- 2'. Antennae not geniculate or without a large obliquely truncate club made up of three segments; or basal margins of elytra armed with ridge or strongly procurved; scutellum small or absent; head may or may not be covered by pronotum; or meso and metatibia unarmed......Not *X. glabratus*
- 3'. Body stout (Fig. 18); or declivity excavated or stongly convex (Figs. 23-28) and without carina on lower lateral margin; tubercles variable but not solely on first and third interstriaNot *X. glabratus*

Citation

Benzel, J. S. 2015. Screening aid: Redbay ambrosia beetle, *Xyleborus glabratus* Eichhoff. Identification Technology Program (ITP), USDA-APHIS-PPQ-S&T, Fort Collins, CO. 7 pp.

References for more information on *Xyleborus glabratus* and non-targets

Anderson, R. S. 2002. Family 131 Curculionidae. pp 722-815. *in*, Arnett R. H. *American Beetles: Volume II.* CRC Press. Boca Raton Florida. 861 pp.

CERIS. 2011. Exotic Wood Borer/ Bark Beetle Survey Reference: *Xyleborus glabratus.* https://caps.ceris. purdue.edu/dmm/1979.

Rabigala, R. J., Dole, S. A., and Cognato, A. I. 2006. Review of American Xyleborina (Coleoptera: Curculeonidae: Scolytinae) Occurring North of Mexico, with an Illustrated Key, *Annuls of the Entomological Society of America*. 99: 1034-1056

Wood, S. L. 1982. The Bark and Ambrosia Beetles of North and Central America (Coleoptera: Scolytidae) a taxonomic monograph. *Great Basin Naturalist Memoirs No* 6. 1359 pp.

Acknowledgments

Funding for this project was provided to J. S. Benzel through section 10201 of the FY2014 Farm Bill. I would like thank Terrence Walters (USDA-APHIS-PPQ-S&T ITP) and USDA-APHIS-PPQ National Identification Services for support of this work and access to imaging equipment. Boris Kondratieff, Donald Bright, and Todd Gilligan (Colorado State University) provided advice on species identification, image editing, and screening aid formatting.