

Lissachatina fulica

Scientific Name

Lissachatina fulica (Bowdich, 1822)

Synonyms:

Achatina fulica Bowdich, 1822

Common Names

Giant African snail, giant east African snail, giant African land snail, escargot Géant

Type of Pest

Mollusk

Taxonomic Position

Class: Gastropoda, **Order:** Pulmonata, **Family:** Achatinidae

Reason for Inclusion

CAPS Target: AHP Prioritized Pest List for FY 2011

Pest Description

From Bequaert (1950):

Eggs: “The egg is broadly ellipsoidal, 5 to 5.5 mm by 4 to 4.5 mm [approx $\frac{3}{16}$ in]... white or slightly yellowish, with a very thin and brittle calcareous shell.”

Adults: “When full-grown, the shell of *L. fulica* consists of from 7 to 9 (very exceptionally 10) whorls, with a moderately swollen body-whorl and a sharply conical spire, which is distinctly narrowed but scarcely drawn out at the apex. The outline varies greatly, even in the same colony, from very slender to moderately obese, the broader specimens tending to be shorter for the same number of whorls. All whorls are decidedly convex, due to the broadly impressed sutures. The aperture is relatively short, even in the broadest specimens, being always shorter than the spire, often



Figure 1. Giant African snail (Yuri Yashin, achatina.ru, Bugwood.org)



Figure 2. Eggs of *Lissachatina fulica* with size reference (David Robinson, USDA-APHIS-PPQ).

considerably so. The outer lip is usually sharp and thin, rarely somewhat thickened or even slightly expanded in very old specimens; it is very convex, evenly curved into a regular semi-ellipse, and inserted on the body-whorl at a sharp, open angle, the upper part of the body-whorl being scarcely or not flattened behind the lip. The columella is more or less concave, sometimes rather weakly so, in which case may be slightly or even much twisted; it tends to be more concave in the broader shells. It should be noted that in *L. fulica* all stages, from the nepionic shell on, have the umbilical slit completely closed and the columella truncate. In all specimens seen, which on general shape and sculpture were referable to *L. fulica*, both columella and parietal callus are white or bluish-white., without any trace of pink.”

The newly hatched, nepionic shell is 5 to 5.5 mm. long and about 4.5 mm wide [approx $\frac{3}{16}$ in], of 2½ nearly smooth whorls, without any granulation or decussation and with only the weakest traces of vertical wrinkles. The first half post-nepionic whorl is more distinctly wrinkled vertically, but not yet decussate; after this the vertical growth striae become gradually stronger and are now cut by spiral engraved lines, into elongate, vertical welts; at first weak, the decussation gains in strength on the fourth and fifth whorls, where it is visible to the naked eye, after which it decreases again, being superficial on the sixth and usually lacking on the seventh and succeeding whorls. In young shells, the periostracum of the early post-nepionic whorls shows in addition to the decussation a superficial, microscopic criss-cross texture, as if a finely woven cloth had been pressed onto the surface; but no trace of this remains on older shells. In the largest full-grown adult shells, of 7 to 9 whorls, the body whorl is nearly even, the growth-striae being very low or superficial, except below the suture where they form short, strong folds, very lightly crenulating the irregular sutural line. When the periostracum is fresh and intact, the terminal whorls appear not only smooth, but also glossy.... When present in the adult, the darker markings appear almost at once on the first post-nepionic whorl as faint, vertical, straight, pale-brown streaks; in very young shells, these streaks stop at the periphery, forming there slightly deflected spots.”

Shell coloration may be variable due to environmental conditions and diet (reviewed in USDA-APHIS, 2007). An identification guide for *L. fulica* can be found in USDA-APHIS (2007). A description of the shell and measurements can be found in Mitra et al. (2005).



Figure 3. *Lissachatina fulica* shell (David Robinson, USDA-APHIS-PPQ).

Biology and Ecology

“This species is highly adaptable to a wide range of environments, modifying its life cycle to suit local conditions. It is one of the greatest threats to agriculture and the environment worldwide due to its reproductive capacity, destructiveness to plants, threat to human health, and large size. [*Liss*] *achatina fulica* prefers environments that are rich in calcium carbonate, such as limestone, marl, and developed areas with an abundance of cement or concrete” (USDA-APHIS, 2007). Optimal development of this species requires a calcium carbonate content of the substratum (calcium in the soil) of at least 3 – 4%. Lack of this results in slow growth, high mortality, cannibalism, and a cease to reproduction (Godan, 1983).

“Snails reach sexual maturity in less than one year. The average life span is 3-5 years, but individuals may reach the age of 9 years” (USDA-APHIS, 2007). *Lissachatina fulica* is a hermaphroditic snail but is required to mate to reproduce. This species uses neither a dart nor a spermatophore when mating (Chase, 2001). Egg laying can begin around six months of age with snails remaining fecund for approximately 400 days (Smith and Fowler, 2003). “Individuals produce from 10 to more than 400 eggs, 8 to 20 days after mating. Under optimum conditions, 300 to 1000 eggs in 3 to 4 batches may be laid each year” (USDA-APHIS, 2007). Hatching occurs after about 11 days. Hatchlings of *L. fulica* will remain beneath the soil surface for a few to several days while consuming their eggshells and organic matter within the soil (Raut and Barker, 2002).

After emergence from the soil, immature larvae begin to feed voraciously; snails ranging from 5 to 30 mm prefer living vegetation while smaller and older snails prefer decaying vegetation and detritus (Smith and Fowler, 2003). Growth rate of this species is correlated with feeding and excretion rate, which is tied to dietary calcium concentrations (reviewed in Dimitriadis, 2001).

Gradually, the smaller snails will disperse from their hatching site and establish home sites. Each day, the snails forage throughout the night and return to their home sites before dawn (Raut and Barker, 2002).

“This species is nocturnal, but may become active at twilight if the day is overcast and the soil is moist and warm. The snail is extremely sensitive to high rates of evaporation. Under moisture stress, it becomes inactive and begins aestivating within 24 hours [aestivation is when an animal becomes dormant during hot and dry weather conditions]. However, aestivation can occur independent of moisture. Scientists believe that aestivation may be cyclic. Snails may aestivate as they cling to objects, aiding in their inadvertent spread to new areas on cargo, vehicles or machinery.

During unfavorable periods, the snail buries itself 10 to 15 cm (4 to 6 inches) deep in soft soil and may become inactive for up to a year, losing 60% of its

weight. Physiological changes in blood and certain organs occur before and during the period of inactivity. This species can reproduce in areas that are too dry for other large snails (Hardouin et al. 1995) (Snail Draft Generic Action Plan, 1986; modified by IICA, 2002; Srivastava, 1985)” (USDA-APHIS, 2007).

Although *L. fulica* is considered a tropical species, it can remain active in temperatures ranging from 9 to 29°C (48.2 to 84.2°F). *Lissachatina fulica* can hibernate to survive temperatures of 2°C (35.6°F) and aestivate in temperatures of 30°C (86°F) (Smith and Fowler, 2003). *Lissachatina fulica* can remain in this state for several months (Smith and Fowler, 2003). Aestivation occurs in moist soil but can occur at sites above-ground (Raut and Barker, 2002).

Damage

Lissachatina fulica prefers to feed on plants in the seedling or nursery stage. Damage by *L. fulica* to the seedling stage can be so severe that farmers change crop species grown. In more mature plants, the type of damage can vary depending on the plant species. Symptoms can range from defoliation to damage of stems, fruits or flowers (Raut and Barker, 2002).

Visual signs of *L. fulica* can include chewing or rasping damage to plants, presence of eggs, juveniles and adults, empty snail shells, mucus and slime trails, large-ribbon like feces, and/or an increase in rat population densities in an area (USDA-APHIS, 2007).

Pest Importance

Lissachatina fulica is considered a significant pest of agricultural crops as well as a vector of a number of human, plant, and animal pathogens. If introduced into the United States, this species may have a high economic impact because of crop damage and the costs of control and



Figure 4. Damage to Leaves of *Dieffenbachia* spp. Caused by *Lissachatina fulica* (Department of Plant Industry, Florida)



Figure 5. Damage to Leaves of *Heliconia* spp. Caused by *Lissachatina fulica* (Department of Plant Industry, Florida).

survey measures (Venette and Larson, 2004). Efforts by the United States to eradicate *L. fulica* from Florida in the late 1960s into the 1970s cost an estimated \$700,000. If the infestation had gone undetected, it is estimated that annual losses would have reached \$11 million (in 1969 dollars) (USDA-APHIS, 1982). Raut and Barker (2002) also state that “opportunity losses associated with enforced changes in agricultural practice, such as limiting the crop species to be grown in a region to those resistant to *L. fulica*” may occur.

This species has been recorded killing young banana plants and destroying the seedlings of *Phaseolus* sp. It also eats the leaves and fruits of plants under cultivation in fields and gardens, including cereals. Damage has been recorded on *Asplenium nidus*, the bread fruit tree (*Artocarpus* sp.), seedlings, flowers, seeds and young pods of cocoa (*Theobroma cacao*), papaya, citrus, young *Hevea* plants, cacti (*Cactus*, *Cereus*, *Opuntia*), the leaves of *Amaranthus* sp., *Bryophyllum* sp., and *Canna* sp., among others (Godan, 1983).

This species has previously threatened production of bananas and coconuts in Western Samoa; displaced small agricultural producers in Brazil, causing an increase in food prices and importation of food; and caused severe vegetable crop damage in parts of India (reviewed in USDA-APHIS, 2007; Cowie, 2006). It is a pest of *Gossypium* (cotton) seedlings in Mauritius, *Artocarpus* (bread fruit tree) in Saipan and various Pacific Islands, *Crotalaria* (East Indian hemp) in Indonesia, *Coffea* (coffee tree) in Tanzania, *Theobroma cacao* (cocoa tree) in Sri Lanka, *Hevea* (rubber tree) in Malaysia and Sri Lanka, *Crotalaria* (rattlebox) in Indonesia, Leguminosae in Malaysia, *Phalaenopsis* (Orchidaceae) in Philippines and Java, and *Asplenium nidus* (bird’s nest fern) in Saipan (reviewed in Godan, 1983). In the tropics, this species is a known pest of vegetables including: cabbage, beans, peas, tobacco, and young plants of *Ipomoea* and *Hevea* (reviewed in Godan, 1983).

Other negative consequences of *L. fulica* establishment could include damage to native plants, alterations to nutrient cycling in the environment, adverse effects to native mollusks (through competition or control measures applied), and effects on human health (Raut and Barker, 2002). When populations are high, the calcium carbonate in the shells can neutralize acid soils, altering the soil properties and affecting the



Figure 6. Damage to Leaves of *Sanchezia nobilis* caused by *Lissachatina fulica* (Department of Plant Industry, Florida).

types of plants that can grow in the soil (reviewed in Cowie, 2006).

Known Food Sources*

This species is polyphagous. The following food source list is from USDA-APHIS (2007). This host list is not all inclusive as reported food sources in the literature tend to focus on geographically important plant species.

Primary Food Sources	Common Names
<i>Albizia lebbbeck</i>	Woman's tongue
<i>Albizia</i> spp.	Albizia
<i>Alsophila</i> spp.	Alsophila
<i>Alstonia scholaris</i>	Devil tree of India
<i>Amaranthus blitum</i>	Purple amaranth
<i>Amaranthus gangeticus</i>	Molten fire
<i>Amaranthus tricolor</i>	Chinese amaranth
<i>Amaranthus viridis</i>	Green amaranth
<i>Amaranthus</i> spp.	Amaranth
<i>Annona muricata</i>	Soursop
<i>Arachis hypogaea</i>	Peanut
<i>Artocarpus altilis</i>	Breadfruit
<i>Artocarpus heterophyllus</i>	Jackfruit
<i>Artocarpus</i> spp.	Breadfruit
<i>Asplenium nidus</i>	Bird's nest fern
<i>Averrhoa bilimbi</i>	Blimb
<i>Basella alba</i>	Ceylon spinach
<i>Basella rubra</i>	Malabar spinach
<i>Blechum brownei</i>	Browne's blechum
<i>Boerhavia diffusa</i>	Boerhavia
<i>Bougainvillea speciabilis</i>	Bougainvillea
<i>Bougainvillea</i> spp.	Bougainvillea
<i>Brassica oleracea</i>	Kohlrabi
<i>Brassica oleracea</i> var. <i>acephala</i>	Cabbage
<i>Brassica oleracea</i> var. <i>botrytis</i>	Cauliflower
<i>Brassica oleracea</i> var. <i>capitata</i>	Cabbage
<i>Brassica oleracea</i> var. <i>italica</i>	Sprouting broccoli
<i>Brassica</i> spp.	Cabbage
<i>Bryophyllum pinnatum</i>	Air plant
<i>Cajanus cajan</i>	Pigeonpea
<i>Calophyllum inophyllum</i>	Indian laurel
<i>Canavalia gladiata</i>	Sword jackbean
<i>Canna edulis</i>	Arrowroot
<i>Canna indica</i>	Canna
<i>Canna</i> spp.	Canna
<i>Carica papaya</i>	Papaya
<i>Cassia</i> (= <i>Chamaecrista</i>) <i>mimosoides</i>	Chamaecrista
<i>Cassia occidentalis</i>	Septic weed
<i>Centrosema pubescens</i>	Flor de conchitas
<i>Cestrum nocturnum</i>	Night Queen

Primary Food Sources	Common Names
<i>Cichorium endivia</i>	Endive
<i>Cichorium intybus</i>	Chicory
<i>Chrysanthemum</i> spp.	Chrysanthemum
<i>Cinnamomum tamala</i>	Indian bay leaf
<i>Citrullus vulgaris</i> (=lanatus)	Watermelon
<i>Citrus limon</i>	Lemon
<i>Citrus reticulata</i>	Tangerine
<i>Citrus sinensis</i>	Sweet Orange
<i>Citrus</i> spp.	Citrus
<i>Clitoria ternatea</i>	Butterfly pea
<i>Colubrina asiatica</i>	Asian nakedwood
<i>Corchorus capsularis</i>	Jute
<i>Corchorus</i> spp.	Corchorus
<i>Cosmos</i> spp.	Cosmos
<i>Crinum</i> spp.	Crinum lily
<i>Crotalaria anagyroides</i>	Crotalaria
<i>Crotalaria pallida</i> var. <i>obovata</i>	Smooth rattlebox
<i>Cucumis melo</i>	Cantaloupe
<i>Cucumis sativus</i>	Cucumber
<i>Cucumis</i> spp.	Melon
<i>Cucurbita maxima</i>	Winter squash
<i>Cucurbita pepo</i>	Field pumpkin
<i>Cucurbita</i> spp.	Gourds
<i>Cyathea lunulata</i>	Tree fern
<i>Daucus carota</i>	Carrot
<i>Dioscorea alata</i>	Greater Yam
<i>Dioscorea bulbifera</i>	Air Yam
<i>Dioscorea</i> spp.	Yam
<i>Dolichos</i> spp.	Beans
<i>Epipremnum aureum</i>	Golden pothos
<i>Erythrina lithosperma</i> (=subumbrans)	Erythrina
<i>Erythrina</i> spp.	Erythrina
<i>Falcataria moluccana</i>	Peacocks plume
<i>Ficus hispida</i>	Fig
<i>Fragaria x ananassa</i>	Strawberry
<i>Gazania rigens</i>	Treasure-flower
<i>Glycine max</i>	Soybean
<i>Glycine</i> spp.	Beans
<i>Gossypium herbaceum</i>	Cotton
<i>Gossypium</i> spp.	Cotton
<i>Grewia mariannensis</i>	
<i>Gynandropis speciosa</i>	The queen's plume
<i>Hevea brasiliensis</i>	Rubber
<i>Hibiscus esculentus</i>	Okra
<i>Hibiscus mutabilis</i>	Dixie rosemallow Land-lily
<i>Hibiscus rosa-sinensis</i>	China rose
<i>Hibiscus</i> spp.	Hibiscus
<i>Hemigraphis colorata</i>	Broad leaf flame ivy

Primary Food Sources	Common Names
<i>Impatiens balsamina</i>	Balsam
<i>Indigofera suffruticosa</i>	Anil de pasto
<i>Ipomoea alba</i>	Tropical white morning-glory
<i>Ipomoea pes-caprae</i>	Beach morning glory
<i>Pachystachys coccinea</i>	Cardinal's guard
<i>Lablab purpureus</i>	Lablab Bean
<i>Lactuca indica</i>	Lettuce
<i>Lactuca sativa</i>	Lettuce
<i>Lactuca</i> spp.	Lettuce
<i>Lagenaria leucantha</i>	Haired gourd
<i>Lagenaria siceraria</i>	Bottle Gourd
<i>Lagenaria vulgaris</i>	Gourd
<i>Lagenaria</i> spp.	Gourds
<i>Laportea crenulata</i>	Tree nettle
<i>Leucaena leucocephala</i>	White leadtree
<i>Catharanthus) rosea</i>	Madagascar periwinkle
<i>Luffa aegyptiaca</i>	Dishcloth gourd
<i>Luffa acutangula</i>	Angled luffa
<i>Luffa cylindrica</i>	Smooth luffa
<i>Luffa</i> spp.	Vegetable sponge
<i>Manihot esculenta</i>	Cassava
<i>Melanolepis multiglandulosa</i>	Alom
<i>Mimosa invisa (=diplotricha)</i>	Giant false sensitive plant
<i>Morinda citrifolia</i>	Indian mulberry
<i>Momordica charantia</i>	Balsam apple
<i>Momordica cochinchinensis</i>	
<i>Momordica</i> spp.	Momordica
<i>Musa acuminata x balbisiana</i>	French plantain
<i>Musa paradisiaca</i>	Plantain
<i>Musa sapientum</i>	Banana
<i>Musa</i> spp.	Bananas
<i>Pachyrhizus erosus</i>	Yam bean
<i>Parkia</i> spp.	Parkia
<i>Passiflora foetida</i>	Fetid passionflower
<i>Passiflora</i> spp.	Passion flower
<i>Pauinia cupana</i>	Guarana
<i>Vigna radiata</i>	Bean
<i>Phaseolus vulgaris</i>	Kidney bean
<i>Physalis peruviana</i>	Peruvian groundcherry
<i>Piper nigrum</i>	Pepper
<i>Pipturus albidus</i>	Waimea pipturus
<i>Pipturus argenteus</i>	Native mulberry
<i>Portulaca grandiflora</i>	Purslane
<i>Portulaca oleracea</i>	Little hogweed
<i>Portulaca</i> spp.	Nine-O'Clock
<i>Prunus persica</i>	Peach
<i>Pueraria montana var. lobata</i>	Kudzu
<i>Raphanus sativus</i>	Radish

Primary Food Sources	Common Names
<i>Ricinus communis</i>	Castor
<i>Sanchezia nobilis</i>	Sanchezia
<i>Scaevola sirecea</i>	Naupaka
<i>Sesamum indicum</i>	Sesame
<i>Spilanthes acmella</i>	Paracress
<i>Spinacea oleracea</i>	Garden spinach
<i>Tabernaemontana divaricata</i>	Pinwheel flower
<i>Tagetes erecta</i>	African marigold
<i>Tagetes patula</i>	Indian marigold
<i>Tagetes</i> spp.	Marigolds
<i>Tectaria</i> spp.	Halberd fern
<i>Tectona grandis</i>	Teakwood
<i>Tephrosia candida</i>	White hoarypea
<i>Tephrosia vogelii</i>	Vogel's tephrosia
<i>Theobroma cacao</i>	Cacao
<i>Thespesia populnea</i>	Portia tree
<i>Trichosanthes dioica</i>	Pointed gourd
<i>Triticum aestivum</i>	Wheat
<i>Vigna radiata</i>	Mung bean
<i>Vigna sinensis unguiculata</i>	Blackeyed pea
<i>Vitis vinifera</i>	Grape
<i>Vigna marina</i>	Notched cowpea
<i>Xanthosoma maffafa</i>	Golden delicious

Secondary Food Sources	Common Name
<i>Abelmoschus esculentus</i>	Okra
<i>Allangana lamarcana</i>	Ballabhi-anga
<i>Allium cepa</i>	Onion
<i>Alocasia indica</i>	Arum
<i>Alocasia macrorrhizos</i>	Giant taro
<i>Aloe indica</i>	Aloe
<i>Amorphophallus campanulatus</i> (= <i>paeoniifolius</i>)	Elephant-Foot Yam
<i>Antigonum leptopus</i>	Antigonon
<i>Alocasia</i> spp.	Elephant ear
<i>Arctium lappa</i>	Greater burdock
<i>Areca catechu</i>	Betel nut palm
<i>Averrhoa carambola</i>	Carambola
<i>Bauhinia accuminata</i>	White dwarf orchard tree
<i>Beta vulgaris</i>	Common beet
<i>Bikkia mariannensis</i>	
<i>Boehmeria nivea</i>	Chinese grass
<i>Brassica campestris</i>	Field mustard
<i>Broussonetia papyrifera</i>	Paper mulberry
<i>Callicarpa cana</i>	
<i>Camellia sinensis</i>	Tea
<i>Capparis cordifolia</i>	Maiapilo
<i>Capsicum annuum</i>	Cayenne pepper

Secondary Food Sources	Common Name
<i>Capsicum baccatum</i>	Locoto
<i>Capsicum</i> spp.	Chili peppers
<i>Senna sophora</i>	Kasuandi
<i>Catharanthus roseus</i>	Periwinkle
<i>Cerbera manghas</i>	Madagascar ordeal bean
<i>Cereus hildmannianus</i>	Hedge cactus
<i>Cereus</i> spp.	Cactus
<i>Cleome gyandra</i>	Spiderwisp
<i>Clerodendron inerme</i>	Glory bower
<i>Coccinia cordifolia</i>	Coccinea
<i>Cocos nucifera</i>	Coconut
<i>Coffea arabica</i>	Arabian coffee
<i>Coffea canephora</i>	Robusta coffee
<i>Coffea</i> spp.	Coffee
<i>Colocasia antiquorum</i>	Arum
<i>Colocasia esculenta</i>	Dasheen or taro
<i>Dalbergia sissoo</i>	Indian rosewood
<i>Dhalia</i> spp.	Dhalias
<i>Dieffenbachia seguine</i>	Dumbcane
<i>Dracaena</i> spp.	Dracaena
<i>Edgaria darjeelingensis</i>	Squash
<i>Elaeis guineensis</i>	Oil palm
<i>Epipremnum pinnatum</i>	Pothos
<i>Eranthemum</i> spp.	Eranthemum
<i>Eucalyptus deglupta</i>	Indonesian gum
<i>Eucalyptus</i> spp.	Australian gum
<i>Eugenia</i> spp.	Star apple
<i>Euphorbia pulcherrima</i>	Poinsettia
<i>Euphorbia trigona</i>	Sandmat
<i>Ficus tinctoria</i>	Fig
<i>Galinsoga parviflora</i>	Gallant-soldier
<i>Gardenia augusta</i>	Gardenia
<i>Gliricidia sepium</i>	Madre de Cacao
<i>Commelina benghalensis</i>	Tropical day flower
<i>Gomphrena globosa</i>	Globe amaranth
<i>Helianthus annuus</i>	Sunflower
<i>Hernandea ovigera</i>	
<i>Ipomoea batatas</i>	Sweet potato
<i>Jasminum sambac</i>	Jasmine
<i>Kalanchoe pinnatum</i>	Kalanchoe
<i>Lycopersicon esculentum</i>	Tomato
<i>Mentha repens</i>	Hortela
<i>Monstera deliciosa</i>	Split leaf philodendron
<i>Montanoa hibiscifolia</i>	Tree daisy
<i>Moringa oleifera</i>	Horseradish tree
<i>Morus alba</i>	White mulberry
<i>Muntingia calabura</i>	Strawberry tree
<i>Nerium</i> spp.	Oleander

Secondary Food Sources	Common Name
<i>Nicotiana</i> spp.	Tobacco
<i>Ochrosia mariannensis</i>	Lipstick tree
<i>Ochrosia oppositifolia</i>	Bwa sousouri
<i>Operculina turpethum</i>	St. Thomas lidpod
<i>Opuntia</i> spp.	Cholla cactus
<i>Pandanus</i> spp.	Screwpine
<i>Pemphis acidula</i>	Small-leafed mangrove
<i>Phalaenopsis</i> spp.	Moth orchids
<i>Pisum sativum</i>	Garden pea
<i>Pisum</i> spp.	Pea
<i>Pluchea indica</i>	Pluchea
<i>Pisidium guaja</i>	Guava
<i>Psychotria mariana</i>	
<i>Tradescantia spathacea</i>	Moses in the cradle
<i>Rosa</i> spp.	Roses
<i>Ruta graveolens</i>	Common rue
<i>Saccharum officinarum</i>	Sugarcane
<i>Salvia</i> spp.	Sage
<i>Sanseveria trifasciata</i>	Snake Plant
<i>Sechium edule</i>	Chayote
<i>Semibarbula orientalis</i>	Moss
<i>Sinapis arvensis</i>	Charlock mustard
<i>Solanum melongena</i>	Eggplant
<i>Solanum tuberosum</i>	Potato
<i>Swietenia mahogoni</i>	Mahogany
<i>Symphytum officinale</i>	Common comfrey
<i>Synedrella nodiflora</i>	Synadrelia
<i>Thea sinensis</i>	Tea
<i>Tradascantia spathacea</i>	Moses in the boat or oysterplant
<i>Trema orientale</i>	Oriental trema
<i>Trichosanthes anguina</i>	Snake gourd
<i>Tridax argentea</i>	Tridax
<i>Vanda</i> spp.	Vanda orchid
<i>Vanilla</i> spp.	Vanilla
<i>Vernonia scandens</i>	Vernonia
<i>Xanthosoma brasiliense</i>	Cakakib
<i>Zea mays</i>	Corn
<i>Zinnia linearis</i>	Zinnia

Many mollusks find food through smell. *Lissachatina fulica* has been found to orient towards the smell of sliced cucumber or carrot (reviewed in Speiser, 2001).

*Terrestrial mollusks do not show host specificity and can feed on multiple crops as well as other materials, like decaying organic matter.

Pathogen or Associated Organisms Vectored

Human and Animal Pathogens

Lissachatina fulica vectors the nematode *Angiostrongylus cantonensis* which can cause eosinophilic meningoenzephalitis in humans as well as *A. costaricensis* which causes abdominal angiostrongylosis (Thiengo et al., 2010). The spread of this disease has been correlated with the spread of *L. fulica* (reviewed in USDA-APHIS, 2007). Achatinids can also carry other diseases that affect humans and animals (Raut and Barker, 2002), including the bacterium *Aeromonas hydrophila* (Cowie, 2006). *Lissachatina fulica* has been experimentally infected with *Anafilaroides rostratus*, a parasite of the bronchii of cats (Malek, 1980).

In Thailand, this considered the most important intermediate host of *Angiostrongylus cantonensis*. It can infect drinking water that *L. fulica* have come in contact with (Godan, 1983).

Note: While most cases of human infections result from consumption of raw or partially cooked snail meat, government inspectors, officers and field surveyors are at-risk due to the handling of live snail, samples, and potential exposure to mucus secretions. **Wear gloves when handling mollusks and wash hands thoroughly after any mollusk survey or inspection activities.**

Plant Pathogens

Lissachatina fulica has been implicated in transmitting *Phytophthora palmivora* (in black pepper, betel pepper, coconut, papaya), *Phytophthora colocasiae* (in taro), and *Phytophthora parasitica* (in eggplant and tangerine); however, the relative importance of *L. fulica* as a transmission agent has not been well established (Raut and Barker, 2002). Plant pathogens are spread through the feces (reviewed in Cowie, 2006).

Known Distribution

Africa: Annobón (Equatorial Guinea), Comoros, Ethiopia, Ghana, Ivory Coast, Kenya, Madagascar, Morocco, Mozambique, Nigeria, Somalia, Seychelles, South Africa, Tanzania (including Zanzibar), and Togo; **Asia:** Andaman Islands, Bangladesh, Bhutan, Bonin Islands, Brunei, Burma, Cambodia, China (including Hong Kong) Taiwan, India, Indonesia, Japan, Laos, Malaysia, Maldives, Nepal, Nicobar Islands, Philippines, Singapore, Sri Lanka, Thailand, and Vietnam; **Caribbean:** Antigua (Antigua and Barbuda), Barbados, Cuba, Guadeloupe, Marie-Galante, Martinique, Saint Lucia, and Trinidad (Trinidad and Tobago); **Oceania:** Caroline Island (Kiribati), Christmas Island (Australia), Cook Islands, Papua New Guinea (including Bismarck Archipelago), Federated States of Micronesia, French Polynesia (including Society Islands), Guam, Hawaii, Mariana Islands, Marshall Islands, Mauritius, New Caledonia, Palau, Réunion, Samoa, Solomon Islands, Tokelau (New Zealand), Tonga, Vanuatu, Wake Island, and Wallis and Futuna; **South America:** Argentina, Bolivia, Brazil, Colombia, Ecuador, and Venezuela (Robinson, 2002; Prasad et al., 2004;

Venette and Larson, 2004; Mitra et al., 2005; NAPPO, 2008; Robinson, 2014, personal communication; Vazquez Perera and Sanchez Noda, 2014).

Pathway

This species has been introduced both purposefully and accidentally into many parts of the world. Intentional movement for medicinal purposes, food (escargot), and research purposes has occurred (reviewed in Cowie, 2006).

“Today, ethnic populations utilizing achatinids as dietary, medicinal, or cultural foodstuffs move live snails with their personal possessions. Most achatinids that PPQ intercepted during the period January 1993 to December 2003 arrived on flights that originated in countries that have established populations of achatinids—countries such as Ghana, Nigeria, and Senegal (Mead 1961, 1979).

Prior to 1997, live [achatinid] snails were seized by PPQ in Arizona, California, Florida and Ohio. During a blitz conducted by Safeguarding, Intervention, and Trade Compliance (SITC) in 2004, PPQ Officers seized and destroyed 6,719 achatinids in six states and 64 cities” (USDA-APHIS, 2007).

This species can also move accidentally on agricultural products (most likely in the egg or juvenile stage). It has previously been intercepted in baggage, cargo, soil (with or without plants), flowers, seeds, and *Zoysia* sp. (peat) (Godan, 1983), among other things.

Potential Distribution within the United States

Lissachatina fulica has been found and subsequently eradicated several times in the United States. This pest was found in the gardens of San Pedro, California in the late 1940s (Abbott, 1949), in Arizona in 1958 (Mead, 1959a) and in Florida in the late 1960s (USDA-APHIS, 2007). In 2004, it was discovered that *L. fulica* was being imported by the pet trade and educational institutions; over 6,700 snails were confiscated from nine states and Puerto Rico to prevent establishment (USDA-APHIS, 2007).

In September 2011, *L. fulica* was detected in Miami, Florida. APHIS, in partnership with the Florida Department of Agriculture and Consumer Services, is currently conducting a regulatory program to eradicate the species. For more information on the efforts in Florida, see the APHIS program page:

http://www.aphis.usda.gov/wps/portal/aphis/ourfocus/planthealth?1dmy&urile=wc m%3apath%3a/aphis_content_library/sa_our_focus/sa_plant_health/sa_domestic_pests_and_diseases/sa_pests_and_diseases/sa_mollusks/ct_giant_african_snail_home.

This species was found in Hawaii in 1936 and was subsequently eradicated through intensive control measures. However, *L. fulica* reestablished later on and is currently present in Hawaii (Godan, 1983).

According to USDA-APHIS (2007) the range of this species is limited by temperature, moisture, and availability of calcium. *Lissachatina fulica* could potentially be found in environments that have a minimum temperature of 1°C (34°F), subtropical rainfall, and available calcium and soil pH of 7.0 – 8.0.

If *L. fulica* were to establish, it would most likely survive in the lower southeastern portion of the United States. Based on host availability and climate, states at increased risk include Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Texas. Portions of California and Arizona also have an increased risk. Smith and Fowler (2003) came up with a similar potential distribution for *L. fulica* in the continental United States. Populations were estimated to extend up to 38° latitude, including most of the southern states, up to Maryland in the east, through Texas to California on the west, and north to Washington and the Pacific Northwest.

Survey

CAPS-Approved Method*:

Visual. Specifics on survey information for *L. fulica* can be found in the New Pest Response Guidelines for Giant African Snails (USDA-APHIS, 2007). See the Introduction to the mollusk manual for specific information on visual surveys.

Survey Site Selection

New introductions of mollusks will likely be related to commerce and human-assisted movement. The habitat and land-use type of each survey site may be variable, ranging from agricultural land, to residential or industrial features. When planning the survey route for a particular site, examine the following microhabitats:

- Near heavily vegetated areas, especially gardens and fields where plants have been damaged by feeding;
- Under rocks, asphalt or cement pieces that are in loose contact with the ground surface;
- Under discarded wooden boards and planks, fallen trees, logs, and branches;
- In damp leaf litter (not wet or soggy), compost piles, and rubbish heaps;
- Under flower pots, planters, rubber mats, tires, and other items in contact with the soil;
- Standing rock walls, cement pilings, broken concrete, and grave markers;
- In gardens and fields where plants have been damaged by feeding snails and slugs; and
- At the base of the plants, under leaves, or in the “heart” of compact plants, such as lettuce or cabbage.

This species can occur in agricultural areas, coastal and wetland areas, disturbed areas, natural and planted forests, riparian zones, scrublands and shrublands, and urban areas (reviewed in Cowie, 2006). It thrives in forest edge, modified forest, and plantation habitats (Raut and Barker, 2002).

Trap Placement

Trapping **cannot** be used alone but can be used to supplement visual surveying. Trapping for mollusks is not species-specific and will attract non-target species, including non-mollusks. Platform or baiting traps can be used to supplement visual inspection. Trap placement can occur in the same areas that visual surveys occur.

Time of year to survey

“Conduct detection surveys on an ongoing basis, with repeated visits at the beginning, during, and—or just after the rainy season. Keep in mind that [*Liss*] *achatina fulica* remains active at a range of 9° – 29°C (48° – 84°F). [*Liss*] *achatina fulica* begins hibernating at 2°C (35°F), and begins aestivation at 30o C (86°F). Plan surveys for early morning and overcast days. Achatinids are active on warm nights, early mornings, and overcast and rainy days” (USDA-APHIS, 2007).

*For the most up-to-date methods for survey and identification, see Approved Methods on the CAPS Resource and Collaboration Site, at <http://caps.ceris.purdue.edu/>.

Key Diagnostics/Identification

CAPS-Approved Method*:

Morphological. Identification should be verified by David Robinson at the USDA APHIS National Malacology Laboratory in Philadelphia, PA.

A key identification feature in *L. fulica* is the columella which ends abruptly; this feature is constant throughout the lifespan of the snail (USDA-APHIS, 2004).

A key to terrestrial mollusks (including *Lissachatina fulica*) is found here: <http://idtools.org/id/mollusc/index.php>.

*For the most up-to-date methods for survey and identification, see Approved Methods on the CAPS Resource and Collaboration Site, at <http://caps.ceris.purdue.edu/>.

Easily Confused Species

This pest may be confused with *Euglandina rosea* or *Orthalicus* spp. both of which are present in the United States. This pest may also be confused with pests not currently present in the United States including *Achatina achatina*, *Archachatina marginata* and *Limicolaria aurora*.

References

- Abbott, R. T. 1949.** March of the Giant African Snail. *Natural History*, 58: 68-71.
- Bequaert, J. C. 1950.** Studies in the Achatinidae, a group of African Snails. *Bulletin of the Museum of Comparative Zoology at Harvard College*, 105: 1-216.
- Chase, R. 2001.** Sensory organs and the nervous system. In G. M. Barker [ed] *The Biology of Terrestrial Molluscs*. CABI International. pp. 179-212.
- Cowie, R. H. 2006.** Global Invasive Species Database: *Achatina fulica*. International Union for Conservation of Nature, Species Survival Commission, Invasive Species Specialist Group. Accessed April 24, 2014 from:
<http://www.issg.org/database/species/ecology.asp?si=64&fr=1&sts=sss>.
- Dimitriadis, V. K. 2001.** Structure and function of the digestive system in Stylommatophora. In G. M. Barker [ed] *The Biology of Terrestrial Molluscs*. CABI International. pp. 237-258..
- Godan, D. 1983.** Pest slugs and snails. Biology and control. Springer-Verlag. Berlin.
- Malek, E. A. 1980.** Snail-Transmitted Parasitic Diseases. Volume II. CRC Press. 324 pp.
- Mead, A. R. 1959.** The appearance of the giant African snail in Arizona. *Proc. Hawaiian Ent. Soc.*, 17(1): 85-86.
- Mitra, S. C., A. Dey, and Ramakrishna. 2005.** Pictorial Handbook. Indian Land Snails. Zoological Survey of India. 344 pp.
- NAPPO. 2008.** Phytosanitary Alert System, *Achatina fulica* (Bowdich) First report of giant African snail (*Achatina fulica*) in Antigua. North American Plant Protection Organization.
- Prasad, G. S., D. R. Singh, S. Senani, and R. P. Medhi. 2004.** Eco-friendly way to keep away pestiferous Giant African snail, *Achatina fulica* Bowdich from nursery beds. *Current Science*, 87(12): 1657-1659.
- Raut, S. K. and G. M. Barker. 2002.** *Achatina fulica* Bowdich and other Achatinidae as pests in tropical agriculture. In G. M. Barker (Ed.), *Molluscs as Crop Pests*, 55-114. New York: CAB International.
- Robinson, D.G. 2002.** *Achatina fulica* Bowdich, 1922. IICA Report on Giant African Snails Workshop. United States Department of Agriculture, Animal and Plant Health Inspection Service. Accessed 24 August, 2010 from:
http://www.aphis.usda.gov/plant_health/plant_pest_info/gas/downloads/achatinafulica.pdf
- Smith, J. W. and G. Fowler. 2003.** Pathway Risk Assessment for Achatinidae with emphasis on the Giant African Land Snail *Achatina fulica* (Bowdich), *Archachatina marginata* (Swainson), and *Limicolaria aurora* (Jay) from the Caribbean and Brazil with comments on the related taxon *Achatina achatina* (Linne) intercepted by PPQ. 1-36. United States Department of Agriculture, Animal and Plant Health Inspection Service, Center for Plant Health Science and Technology, Plant Epidemiology and Risk Analysis Laboratory.
- Speiser, B. 2001.** Food and Feeding Behavior. In G. M. Barker (ed.). *The Biology of Terrestrial Molluscs*. CABI International. pp. 259-288.
- Thiengo, S. C., A. Maldonado, E. M. Mota, E. J. L. Torres, R. Caldeira, O. S. Carvalho, A. P. M Oliveira, R. O. Simões, M. A. Fernandez and R. M. Lanfredi. 2010.** The giant African snail

Achatina fulica as natural intermediate host of *Angiostrongylus cantonensis* in Pernambuco, northeast Brazil. *Acta Tropica*, 115: 194-199.

USDA-APHIS. 1982. Pests not known to occur in the United States or of limited distribution, No. 22: giant African snail. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine.

USDA-APHIS. 2004. Giant African Snail *Achatina fulica*. Mollusk Action Plan Work Group. United States Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine.

USDA-APHIS. 2007. New Pest Response Guidelines. Giant African Snails: Snail Pests in the Family Achatinidae. USDA-APHIS-PPQ-Emergency and Domestic Programs-Emergency Planning, Riverdale, Maryland. http://www.aphis.usda.gov/import_export/plants/manuals/

Vazquez Perera, A. A. and J. Sanchez Noda. 2014. Introducción del caracol gigante Africano, vector de meningoencefalitis eosinofílica en una zona urbana de La Habana. Laboratorio de Malacología, Instituto de Medicina Tropical Pedro Kourí. Last accessed August 15, 2014, from <http://instituciones.sld.cu/ipk/introduccion-del-caracol-gigante-africano-vector-de-meningoencefalitis-eosinofilica-en-una-zona-urbana-de-la-habana/>.

Venette, R. C. and M. Larson. 2004. Mini Risk Assessment: Giant African Snail, *Achatina fulica* Bowdich [Gastropoda: Achatinidae]. Cooperative Agricultural Pest Survey Program.

This datasheet was developed by USDA-APHIS-PPQ-CPHST staff. Cite this document as:

Molet, T. 2011. CPHST Pest Datasheet for *Lissachatina fulica*. USDA-APHIS-PPQ-CPHST

Reviewed by: David Robinson, USDA-APHIS National Malacologist.

Revisions

April 2014

- 1) Revised **Biology and Ecology** section
- 2) Revised **Pest Importance** section
- 3) Revised **Pathogen or Associated Organisms Vectored** section
- 4) Added **Pathway** section
- 5) Revised **Potential Distribution within the United States** section
- 6) Revised **Survey** section

August 2014

- 1) Revised **Known Distribution** section
- 2) Revised **Key Diagnostics/Identification** section

August 2016

- 1) NAPPFAST map removed